arXiv:1303.4357v1 [quant-ph] 18 Mar 2013

Quantum Correlations are Tight Bounded by Exclusivity Principle

Bin Yan*
Department of Physics, Purdue University, West Lafayette, Indiana 47906, USA

(Dated: 11 March 2013)

It is a fundamental problem in physics that what principle limits the correlations as predicted by our current description of nature, based on quantum mechanics. One possible explanation is the "global exclusivity" principle recently been discussed in [Phys. Rev. Lett. **110**, 060402 (2013)]. In this work we show that this principle actually has a much stronger restriction on the probability distribution. We provide a tight constraint inequality imposed by this principle and prove that this principle singles out quantum correlations in scenarios represented by any graph. Our result implies that exclusivity principle might be one of the fundamental principles of nature.

PACS numbers: 03.65.Ta, 02.10.Ox, 03.65.Ud

Introduction.—Quantum correlations between observables are contextual and nonlocal such that quantum mechanics(QM) is incompatible neither with local hidden variable (LHV) theories [1] nor with noncontextual hidden variable (NCHV) theories [2–4]. Noncontextuality (NC) inequalities [5–9] and Bell inequalities [10–13] (which are special type of NC inequalities that require spacelike separated tests) are the basic tools to characterize quantum correlations. Intriguingly, the maximum quantum violation of these correlation inequalities are bounded in a very special way. A fundamental open problem is that what is the physical principe that prevents QM to be more contextual [14, 15] or more nonlocal [16–19]?

Recent approaches to the understanding of quantum correlations address this problem into a general probability theory framework [20–23, 26]. The main idea is that every correlation inequality can be associated to a group of events, whose relationships can be represented by a graph G. G is such a graph that each of its vertex represents an event and two vertices are adjacent if the corresponding events are mutually exclusive. Correlations between observables can thus be characterized by the sum S of the probabilities of these events.

It has been shown [21, 23] that for a given set of events $\{u_i\}$ with the corresponding graph G, the maximum value of S for NCHV theories is the *independent number* of G, $\alpha(G)$. While the upper bound for S predicted by QM is given by the *Lovász number*[27] of G:

$$\vartheta(G) = Max\sum_i |\langle\varphi|v_i\rangle|^2, \qquad (1)$$

where $|\varphi\rangle$ and $|v_i\rangle$ are unit vectors in Eucledian space and the maximum is taken in any dimensions over all possible $|\varphi\rangle$ and *orthogonal representation* $\{|v_i\rangle\}$ (which means that each vector $|v_i\rangle$ is assigned to a vertex of G and two vectors are orthogonal if their corresponding vertices are adjacent). $\{|\langle\varphi|v_i\rangle|^2\}$ are QM allowed probability distributions. Now the question is that: Why does QM stop at the upper bound $\vartheta(G)$? The mean idea is to identify "natural" information principles, formulated only with the constraints on probability distribution, that prevent stronger correlations than QM.

One possible explanation is from the exclusivity principle(EP) [21, 23]: the sum of probabilities of pairwise exclusive events cannot exceed 1. Note that in general probability framework, Bool's axiom [28] only demands that the sum of the probabilities of jointly exclusive events cannot exceed 1, while pairwise exclusive events are not necessary jointly exclusive. Thus this principle indeed imposes a nontrivial restriction on the potability distributions. This simple principle, originally follows from Specker's conjecture on the basic principle of QM(see Ref. [25] for a survey), has been used recently [21, 23, 26] to investigate its fundamental role in bounding the quantum correlations.

Denote $\{P_i\}$ as the probabilities for a given set of events. Clearly, according to this principle, the sum of probabilities of any pairwise exclusive events in $\{u_i\}$ can not exceed 1. Thus $\{P_i\}$ should at least satisfy:

$$\sum_{i\in C} P_i \leq 1, \qquad (2)$$

where C is any such subset of $\{u_i\}$ that events in C are pairwise exclusive. Under this constraint, the maximum value of S is the so-called *fractional packing number* of the corresponding graph G. It has been shown [21] that constraint in inequality (2) singles out quantum corrections for a class of scenarios represented by graphs with their fractional packing numbers equal to their Lovász numbers. These scenarios include Bell inequalities for Greenberger-Horne-Zeilinger states [10] and graph states[11–13], and some bipartite Bell inequalities [29, 30] as well as all the state-indepented NC inequalities in Ref. [7]. In a more recent work [23], by applying exclusivity principle to two copies of the Klyachko-Can-Binicioglu-Shumovsky (KCBS) [6] experiments, Cabello showed that the upper bound of KCBS inequality for QM is exactly the maximum value allowed by EP. However, the answer was leaded to an open question when applying this principle to multiple copies of Clauser-Horne-Shimony-Holt(CHSH) experiments. It is still not clear whether the maximum quantum violation of CHSH inequality is tight bounded by EP, or more generally, whether EP can single out quantum correlations in scenarios represented by any graph.

In the following we show that, without any additional assumptions, EP actually has a much stronger restriction on $\{P_i\}$ than inequatlity (2). We provide a tight constraint inequality on $\{P_i\}$ and show that this principle indeed singles out nature's maximum contextuality and nonlcality for any graph.

Probability distributions allowed by EP.—Given a set of events $\{u_i\}$ and the corresponding graph G, we now consider another group of events $\{v_i\}$ which are completely independent with $\{u_i\}$. Events $\{v_i\}$ have the relationships that v_i and v_j are mutually exclusive if u_i is not exclusive with u_j. Therefore the corresponding graph of events $\{v_i\}$ is exactly the complementary graph of G (Graphs depicted in Fig. (1) are examples of complementary graphs). We now consider u_i and v_i together as a joint event u_iv_i (which means event u_iv_i happens if and only if u_i and v_i both happen). Clearly, events $\{u_iv_i\}$ form a pairwise exclusive events set. Donate $\{P_i\}$ and $\{P'_i\}$ the EP allowed probability distributions of $\{u_i\}$ and $\{v_i\}$, respectively. As u_i and v_i are completely independent, the joint probability for event u_iv_i will be $P_iP'_i$.

Note that the joint events $\{u_iv_i\}$ are still real events, despite they are constructed by jointly viewing two set of independent events. A physical principle should be universal such that the probability distributions of events $\{u_iv_i\}$ must be restricted by exclusivity principle:

$$\sum_i P_i {P'}_i \leq 1. \tag{3}$$

Thus the constraint on $\{P_i\}$ is that: $\{P_i\}$ must satisfy inequality (3) for *any* EP allowed probability distribution $\{P'_i\}$ on $\overline{G}$. It is a much stronger restriction imposed by EP since all possible solutions $\{P_i\}$ for this inequality automatically satisfy constraint (2). The above inequality is one of the main results in this work. We remark here that the "global" property assumed in Ref. [23] actually comes from the universality of a physical principle. Instead of imposing restriction only on the probability distributions on the given graph G under consideration, as in inequality (2), exclusivity principle exerts constrains upon the whole probability distribution set including all graphs. It demands the *compatibility* among different graphs. In other words, EP allowed probability distributions are such that they can not generate joint probability distributions which are not allowed by EP. It is in this sense we suggest to call this principle as "Consistent Exclusivity" (CE). Similar proposal, while under different considerations, has also been discussed in [24, 25].

The reason why applying constraint in inequality (2) to two copies of KCBS experiments, after assuming global exclusivity in Ref. [23], singles out quantum contextuality can be explained as this: The graph corresponding to KCBS inequality is a pentagon, whose complementary graph is exactly itself. The OR product of two pentagons is a 25-vertex graph which contains five 5-vertex complete graphs. Apply inequality (2) to each of the five complete graphs would cover the constraint in inequality (3). Actually, as a pentagon is self-complemental, inequality (3) is reduced to a self-constraint inequality $\sum P^2 \leq 1$, which immediately gives us the maximum value of S.

In general cases, calculating the upper bound for the sum of P_i comes down to a linear optimization problem subject to constraint in inequality (3). Now the only problem is that so far we do not know what is exactly the possible probability distribution $\{P'_i\}$ on $\overline{G}$. In fact, EP allowed distributions $\{P'_i\}$ are also bounded by inequality (3) ranging over all EP allowed $\{P_i\}$. This "global" property, that all probability distributions should compatible with each other, makes it difficult to estimate the maximum value of S using inequality (3). Our approach for this problem is to find a subset of all EP allowed $\{P'_i\}$ on $\overline{G}$.

Consider now the QM allowed probability distributions on the given graph G and $\overline{G}$, which (in orthogonal representation) can be written as $\{|\langle\psi|u_i\rangle|^2\}$ and $\{|\langle\varphi|v_i\rangle|^2\}$, respectively. We have the following inequality:

$$\sum_i |\langle\psi|u_i\rangle|^2 |\langle\varphi|v_i\rangle|^2 \leq 1. \tag{4}$$

This is due to the fact that $\{|u_i\rangle \otimes |v_i\rangle\}$ form an orthogonal system such that $\sum_i |\langle\psi|u_i\rangle|^2|\langle\varphi|v_i\rangle|^2 = \sum_i |(\langle\psi| \otimes \langle\varphi|)(|u_i\rangle \otimes |v_i\rangle)|^2$ is always less than 1 for any unit vectors $|\psi\rangle \otimes |\varphi\rangle$.

The meaning of inequality (4) is that all QM allowed probability distributions on graph G and $\overline{G}$ automatically satisfy the restriction imposed by EP in inequality (3). Generally, this fact can be seen from a simple property of ϑ function of any complete graph $G_{complete}$ (in which any two vertices of are adjacent):

$$\vartheta(G_{complete}) = 1, \tag{5}$$

which means that in QM, the sum of probabilities of any group of pairwise exclusive evens can not exceed 1. Namely, for any given graphs, probability distributions allowed by QM constitute the subset of probability distributions allowed by EP.

Note that since EP might allow probability distributions that can not be realized in the orthogonal representation, it is nontrivial to say that maximum quantum violation is exactly the upper bound imposed by EP. In other words, we can not use the orthogonal representation on graph G to estimate the upper bound of the sum of P_i. However, this feather provides a group of possible solutions of $\{P'_i\}$, that is, all QM allowed probability distribution $|\langle\varphi|v_i\rangle|^2$ on $\overline{G}$. $\{P_i\}$ should at least satisfy inequality (3) when $\{P'_i\}$ adopts these distributions.



FIG. 1. (a) Graph G of the relationships of the 8 events $\{u_i\}$ involved in CHSH inequality. (b) Graph of the relationships of the 8 events $\{v_i\}$ completely independent with events $\{u_i\}$. Their relationships form the complementary graph of G. This graph is the 8-vertex (1,2)-circulant graph $Ci_8(1,2)$. Joint events $\{u_iv_i\}$ generate the 8-vertex complete graph.

By ranging $\{P'_i\}$ over all unit vector $|\varphi\rangle$ and orthogonal representation $\{|v_i\rangle\}$ on $\overline{G}$, we now get a relatively weaker constraint for $\{P_i\}$:

$$\sum_i P_i|\langle\varphi|v_i\rangle|^2 \leq 1. \tag{6}$$

This constraint actually, as we will see, is sufficient to single out quantum correlations in any graph.

Maximum quantum violation of CHSH inequality.—We now illustrate how EP singles out the maximum quantum violation for CHSH inequality [31].

The bipartite scenario corresponding to CHSH inequality involves eight events [23] with their relationships being represented by the graph G depicted in Fig. 1(a). The upper bounds of CHSH inequality imposed by noncontextual local hidden variable (NCLHV) theory, QM (the bound is known as Tsireson's bound [32]) and nonsignaling(NS) [16] can be expressed as the following:

$$\sum P_i \leq 3^{\ NCLHV} \leq 2+\sqrt{2}^{\ QM} \leq 4^{\ NS} \tag{7}$$

where the sum is extended to all the 8 events. Many efforts have been made [17–19, 33] to explain why QM stops at Tsireson's bound despite more general probability distributions from a PR box [16] does not violate nonsignaling. In Ref. [23], Cabello showed that the "global exclusivity" rules out nonsignaling with a lower bound than 4, which is $\frac{8}{\sqrt{5}} \approx 3.5778$. (The same value was also obtained in [26] by assuming "local orthogonality", which is essentially the same as exclusivity principle).

Here we use the constraint in inequality (6) and range all possible orthogonal representations on $Ci_8(1,2)$, the complementary graph of G, as depicted in Fig. 1(b). Due to the symmetry of G, we should expect that the CHSH inequality reaches its maximum value with the eight evens being assigned the same probability P, which is bounded by

$$P\sum |\langle\varphi|v_i\rangle|^2 \leq 1. \tag{8}$$

This must be satisfied for any orthogonal representations such that:

$$P \leq \frac{1}{\max\sum\limits_i |\langle\varphi|v_i\rangle|^2} = \frac{1}{\vartheta(\overline{G})}, \tag{9}$$

where the maximum ranges over all orthogonal representations on graph $Ci_8(1,2)$ and $\vartheta(\overline{G}) = \max\sum\limits_i |\langle\varphi|v_i\rangle|^2$ is the well known Lovász number for $Ci_8(1,2)$, which is $8-4\sqrt{2}$. We immidiately get the upper bound for CHSH inequality imposed by EP: $8P_{max} = 2+\sqrt{2}$, which is exactly equal to the maximum quantum violation. It is interesting while quite reasonable to see that this number can be given by the complemental graph of the original graph of the 8 events. That is the result of the "global" property of exclusivity principle.

Generalization of our result—We now give a simple proof to show that for correlation inequalities represented by *any* graph, the upper bounds given by exclusivity principle are exactly the same as QM predicted, such that quantum correlations are tight bounded by exclusivity principle.

As discussed above, probability distributions $\{P_i\}$ on a given graph G are restricted by inequality (6) ranging over all orthonormal representations on $\overline{G}$, the complement graph of G. For a given normalized vector $|\varphi\rangle$ and orthonormal representation $\{|v_i\rangle\}$, we pick out the minimum value among $|\langle\varphi|v_i\rangle|^2$, which satisfies

$$(\sum_i P_i)\min|\langle\varphi|v_i\rangle|^2 \leq \sum_i P_i|\langle\varphi|v_i\rangle|^2 \leq 1, \tag{10}$$

or

$$\sum_i P_i \leq \max\frac{1}{|\langle\varphi|v_i\rangle|^2}. \tag{11}$$

Inequality (11) should hold for any normalized vector $|\varphi\rangle$ and orthonormal representation $\{|v_i\rangle\}$ on graph $\overline{G}$. This further gives us:

$$\sum_i P_i \leq \min\max\frac{1}{|\langle\varphi|v_i\rangle|^2}, \tag{12}$$

where the maximum is taken over the given $\{v_i\}$ and $|\varphi\rangle$ and the minimum ranges over all orthonormal representation $\{|v_i\rangle\}$ and unit vector $|\varphi\rangle$.

The right side of inequality (12) equals to the Lovász function for graph G (see *Lemma* 1 in Ref. [27]), which is exactly the maximum value for QM. The equality in (12) can hold since the orthogonal representation of graph G is EP allowed distribution of $\{P_i\}$, which can achieve the maximum value $\vartheta(G)$. Interestingly, the tight bound is given by only ranging $\{P'_i\}$ over the subset of all EP allowed distribution. This tells us other possible distributions which are not included in the orthogonal representation, if there are any, will not give even lower bound than $\vartheta(G)$.

Conclusion and conjecture.— In this work, we have provided further understanding of the "global exclusivity" principle recently discussed in Ref. [23]. We show that this principle actually has a much stronger restriction on the probability distribution. Instead of imposing constraint only on the given graph under consideration, this principle imposes restriction on the whole probability structure including *all* graphs. We have provided a constrain inequality and show that this principle indeed singles out quantum correlations represented by any graph. Namely, quantum correlations are tight bounded by this simple principle.

It is still not clear whether probability distributions allowed by QM are all that allowed by EP. What we can conclude here is that even if the probability distribution set is wider than that allowed by QM, it will not provide stronger restriction on the correlations. It is in this sense we conjecture that exclusivity principle not only limits quantum correlation, but also at least "almost" determines the probability structure of quantum mechanics. It is remarkable that our result implies this principle, along with other physical principles such as uncertainty and nonsignaling, might be one of the fundamental principles of nature.

---

[*] yanbin@purdue.edu

[1] J. S. Bell, Physics **1**, 195 (1964).
[2] E. P. Specker, Dialectica **14**, 239 (1960).
[3] S. Kochen and E. P. Specker, J. Math. Mech. **17**, 59 (1967).
[4] J. S. Bell, Rev. Mod. Phys. **38**, 447 (1966).
[5] A. Cabello, S. Filipp, H. Rauch, and Y. Hasegawa, Phys. Rev. Lett. **100**, 130404 (2008).
[6] A. A. Klyachko, M. A. Can, S. Binicioğlu, and A. S. Shumovsky, Phys. Rev. Lett. **101**, 020403 (2008).
[7] A. Cabello, Phys. Rev. Lett. **101**, 210401 (2008).
[8] R. W. Spekkens, D. H. Buzacott, A. J. Keehn, B. Toner, and G. J. Pryde, Phys. Rev. Lett. **102**, 010401 (2009).
[9] P. Badziąg, I. Bengtsson, A. Cabello, and I. Pitowsky, Phys. Rev. Lett. **103**, 050401 (2009).
[10] N. D. Mermin, Phys. Rev. Lett. **65**, 1838 (1990).
[11] O. Gühne, G. Tóth, P. Hyllus, and H. J. Briegel, Phys. Rev. Lett. **95**, 120405 (2005).
[12] A. Cabello, O. Gühne, and D. Rodríguez, Phys. Rev. A **77**, 062106 (2008).
[13] A. Cabello, M. G. Parker, G. Scarpa, and S. Severini, arXiv:1211.4250.
[14] A. Cabello, Nature (London) **474**, 456 (2011).
[15] A. Cabello, in *A Computable Universe*, edited by H. Zenil (World Scientific, Singapore, 2012), Chap. 31.
[16] S. Popescu and D. Rohrlich, Found. Phys. **24**, 379 (1994).
[17] M. Pawłowski, T. Paterek, D. Kaszlikowski, V. Scarani, A. Winter, and M. Żukowski, Nature (London) **461**, 1101 (2009).
[18] M. Navascués and H. Wunderlich, Proc. R. Soc. A **466**, 881 (2009).
[19] J. Oppenheim and S. Wehner, Science **330**, 1072 (2010).
[20] J. Barrett, Phys. Rev. A **75**, 032304 (2007)
[21] A. Cabello, S. Severini, and A. Winter, arXiv:1010.2163.
[22] R. Gallego, L. E. Würflinger, A. Acín, and M. Navascués, Phys. Rev. Lett. **107**, 210403 (2011).
[23] A. Cabello, Phys. Rev. Lett. **110**, 060402 (2013)
[24] J. Henson, arXiv:1210.5978
[25] A. Cabello, arXiv:1012.1756
[26] T. Fritz, A. B. Sainz, R. Augusiak, J. Bohr Brask, R. Chaves, A. Leverrier, and A. Acín, arXiv:1210.3018
[27] L. Lovász, IEEE Trans. Inf. Theory **25**, 1 (1979).
[28] G. Boole, Phil. Trans. R. Soc. Lond. **152**, 225 (1862), see p. 230.
[29] A. Cabello, Phys. Rev. Lett. **87**, 010403 (2001).
[30] L. Aolita, R. Gallego, A. Acín, A. Chiuri, G. Vallone, P. Mataloni, and A. Cabello, Phys. Rev. A **85**, 032107 (2012).
[31] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
[32] B. S. Cirel'son [Tsirelson], Lett. Math. Phys. **4**, 93 (1980).
[33] C. Pfister and S. Wehner, arXiv:1210.0194.